



**ENSP**  
European Network  
for Smoking and Tobacco Prevention



**next<sup>+</sup>**  
**ENSP**  
Fighting Tobacco Use  
in Europe

## OPEN LETTER

# Youth Nicotine Addiction: From Prevention to Protection No More Delays, Time to Act Together!

*Tobacco use is unlike other threats to global health. Infectious diseases do not employ multinational public relations firms. There are no front groups to promote the spread of cholera. Mosquitoes have no lobbyists[1].*

To Her Excellency,  
Ursula von der Leyen  
President of the European Commission  
Copy to The College of Commissioners

Dear President von der Leyen,

More than four centuries ago, tobacco was described as “loathsome to the eye, hateful to the nose, harmful to the brain and dangerous to the lungs.” Today, scientific evidence is unequivocal: **tobacco consumption remains the leading cause of preventable death worldwide. Around 27% of all cancers are attributable to tobacco use, and more than seven million people die every year from tobacco-related diseases.**

At the heart of tobacco consumption lies a powerful and often underestimated driver: nicotine addiction. It is this addictive mechanism that sustains use over time and makes cessation difficult. **For children and young people, nicotine represents the primary entry point into long-term dependency.** Often introduced through products that appear attractive, modern, or less harmful, many young users are drawn in without fully understanding the risks, only to find themselves quickly dependent. Addressing nicotine addiction at its source is therefore essential to preventing a new generation from being drawn into lifelong use.

**The European Union has set an ambitious and commendable objective: to create a tobacco-free generation by 2040, as outlined in Europe’s Beating Cancer Plan.** Achieving this goal requires a clear and sustained focus on protecting young people from developing nicotine dependence, which remains the primary pathway into long-term tobacco and nicotine use.

**Across the Union, new nicotine products are rapidly expanding in visibility and accessibility,** often deliberately designed and marketed to appeal to younger audiences through flavours, sleek packaging, and targeted digital promotion. This trend poses a serious and growing public health threat.

[1] <https://pmc.ncbi.nlm.nih.gov/articles/PMC8519313/>

Preventing early exposure to nicotine must be treated as an urgent priority, as addiction developed during adolescence significantly increases the likelihood of lifelong dependence and long-term harm. Crucially, youth nicotine addiction is not an isolated issue. It frequently acts as a gateway to broader patterns of substance use, with strong evidence linking early nicotine exposure to an increased risk of subsequent use of alcohol and illicit drugs.

**At the same time, the health consequences of tobacco and nicotine use are profound, systemic, and too often underestimated.** Tobacco use is a leading cause of cardiovascular disease, including myocardial infarction, arteriosclerosis, cerebrovascular accidents, and peripheral vascular disease, which can result in limb amputations. It is also responsible for a wide range of cancers, extending far beyond lung cancer to include cancers of the bladder, kidney, colorectum, and the pharyngeal, laryngeal, and oral regions. **Framing tobacco-related harm primarily as a respiratory issue fails to capture its full impact on the human body.** These realities underscore the urgent need for comprehensive, forward-looking policies that effectively reduce affordability, limit exposure, and protect young people from nicotine addiction in all its forms.

**Tobacco and nicotine control is not solely a public health issue; it is a complex, cross-sectoral challenge that affects fiscal policy, trade, environment, agriculture, and digital regulation.** The World Health Organization Framework Convention on Tobacco Control (WHO FCTC), ratified by the European Union, provides a comprehensive international framework demonstrating how to address this complexity through coordinated, multisectoral action.

**A healthy and resilient European Union depends on the well-being of its younger generations. A child addicted to nicotine at 13 is far more likely to become a chronically ill adult by 40.**

Reducing nicotine uptake among youth contributes directly to stronger future workforce participation, improved productivity, and more sustainable healthcare systems. **Protecting young people today is therefore closely linked to Europe's long-term competitiveness and social cohesion.**

In this context, a forward-looking regulatory approach is essential. **The current Tobacco Products Directive provides an important foundation**, yet the evolving landscape of nicotine products, including synthetic nicotine and nicotine analogues, calls for a more comprehensive framework. **Revising this directive with the aim of developing a Tobacco and Nicotine Products Directive would allow for a coherent and consistent regulation of all nicotine-containing products.** Such an approach would strengthen the protection of young people by **ensuring that all products with addictive potential are addressed** under a unified public health strategy, reducing regulatory gaps and supporting more effective prevention of youth nicotine initiation.

As it has done over the past 30 years, **the European Network for Smoking and Tobacco Prevention (ENSP) stands ready to support the European Commission in advancing tobacco and nicotine control across the Union.** Established in 1996–1997 at the request of the European Commission to provide expertise in this field, ENSP is not a lobbying organisation.

**Protecting the health of European citizens is not a lobbying activity, but a public interest responsibility.**

ENSP operates as an independent technical partner[2], drawing on its extensive European network, longstanding collaboration with EU institutions and the World Health Organization, and recognised expertise in policy, research, and implementation. **In this capacity, ENSP can contribute to enhanced coordination** among stakeholders, the provision of **evidence-based input**, support for the **implementation of EU legislation** and WHO FCTC provisions, **capacity-building** across Member States, and strengthened **prevention efforts**, particularly among youth.

As President of the European Commission and medical doctor, you continue to shape policies that influence the well-being of millions of Europeans. Strengthening efforts to prevent nicotine addiction among young people represents a powerful opportunity to advance both public health and the future prosperity of the Union.

**Reaching the objective of reducing tobacco use to below 5% by 2040**, as specified also in the ENSP Statutes from 2011, will depend on sustained commitment, coordinated action, and a strong emphasis on prevention. **By prioritizing youth protection and addressing nicotine dependence at its source, the European Union can make meaningful progress** toward a healthier, more resilient future.

The years ahead offer an important opportunity to strengthen this vision and ensure that the next generation grows up less exposed to addiction and better protected in their health and well-being.

Through joint efforts and cooperation, we aim to ensure that the voice of the European people is heard and reflected in our collective action.

**ENSP President:**

Prof. Panagiotis Behrakis - GREECE

**ENSP Board Members:**

Danielle van Kalmthout - BELGIUM

Raquel Fernandez Megina - SPAIN

Prof. Maria Sofia Cattaruzza - ITALY

Helen Stjerna - SWEDEN

Prof. Luke Clancy - IRELAND

Prof. Loïc Josseran - FRANCE

**ENSP Secretariat:**

Cornel RADU LOGHIN - Secretary General

Charis GIRVALAKI - Scientific Director

Karina MOCANU - ENSPNext (Youth Group)  
Coordinator

**ENSP Scientific Committee Coordinator:**

Dr. Pierre BIZEL - BELGIUM

## Signed and supported by:



1. Youth Health Organization International (YHO).
2. AllCan International
3. Global Network for Tobacco-Free Healthcare Services (GNTH).
4. EuroHealthNet
5. Smoke Free Partnership (SFP).
6. European Heart Network (EHN).
7. European Respiratory Society (ERS).
8. Cancer Patients Europe (CPE).
9. European Association of Urology.
10. Lung Cancer Europe
11. Digestive Cancers Europe (DiCE).
12. Pancreatic Cancer Europe
13. European Liver Patients' Association
14. European Academy of Allergy and Clinical Immunology (EAACI).
15. TB Europe Coalition
16. European Kidney Health Alliance (EKHA).
17. Association of European Cancer Leagues (ECL).

## Signed and supported by:



### ALBANIA

Faculty of Health at The Ismail Qemali University of Vlorë)



### AUSTRIA

Austrian Council on Smoking & Health  
VIVID – Institute for the Prevention of Addiction)



### BELGIUM

The Belgian Alliance for a Smoke-Free Society  
Kom op tegen Kanker  
Fondation Contre le Cancer  
Respiratoire Gezondheid VRGT  
OXY Belgium



### BOSNIA & HERZEGOVINA

PROI – Association for Progressive Initiatives



### BULGARIA

Tobacco & Nicotine Free Life Coalition  
Bulgarian Joint Cancer Network (BJCN))



### CZECH REPUBLIC

Cancer Patients Voice



### CYPRUS

Cyprus National Addictions Authority



### DENMARK

Danish Cancer Society  
Danish Heart Foundation  
The Cancer Association Tidslerne



### FINLAND

Suomen ASH ry/ASH Finland  
Syöpäjärjestöt - Cancer Society of Finland



### FRANCE

CONTRE-FEU - L'Alliance contre l'industrie du tabac  
DNF - Demain sera non-fumeur



### GEORGIA

FCTC Implementation and Monitoring Center in Georgia



### GERMANY

Das Aktionsbündnis Nichtraucher e.V. (ABNR)  
German Respiratory Society  
German Network for Tobacco Free Healthcare Services )  
BLUE 21 e.V. / Unfairtobacco



### GREECE

Hellenic Cancer Society  
Hellenic Thoracic Society



### IRELAND

TobaccoFree Research Institute  
The Irish Heart Foundation  
ASH Ireland, Council of the Irish Heart Foundation  
Pancreatic Cancer Ireland



### ITALY

L'Alleanza per un'Italia Senza Tabacco  
Società Italiana di Tabaccologia - SITAB  
Istituto di Ricerche Farmacologiche Mario Negri



### KOSOVO

KADC - Kosovo Advocacy Development Center



## LITHUANIA

Lithuanian Tobacco and Alcohol Control Coalition



## LUXEMBOURG

Fondation Cancer



## MALTA

MAPHM-Malta Association of Public Health Medicine



## MOLDOVA

The Center for Health Policies and Studies (PAS Center)  
Coalition for Respiratory Health  
Association of Oncology Patients "Viața Continuă"



## THE NETHERLANDS

Gezondheidsfondsen voor Rookvrij (Hartstichting, KWF en Longfonds)  
Trimbos Institute



## NORTH MACEDONIA

Macedonian Respiratory Society



## NORWAY

Norwegian Cancer Society



## POLAND

Maria Skłodowska-Curie National Research Institute of Oncology  
Fundacja „Smart Health – Zdrowie w 3D”



## PORTUGAL

SHINE 2Europe



## ROMANIA

Romanian Society of Pneumology  
Healthy Romania Generation 2035 Association / Tobacco-Free Romania Initiative  
The Bucharest University of Economic Studies (ASE), Research Center in International Economic Relations  
AER PUR Romania  
M.A.M.E Association



## SERBIA

Institute of Economic Sciences  
National Initiative of Non-Smokers of Serbia



## SLOVENIA

Slovenian Coalition for Public Health, Environment and Tobacco Control  
No Excuse Slovenia



## SPAIN

Comité Nacional para la Prevención del Tabaquismo - CNPT  
Asociación Española Contra el Cáncer  
Institut Català d'Oncologia  
La Sociedad Española de Medicina de Familia y Comunitaria - semFYC  
El Colegio Oficial de Dentistas de Las Palmas (COELP) - Fundación Canaria Dental Nofumadores  
Sepa – Fundación Sepa  
ATAEX - Asociacion tabaquismo de Extremadura  
APTA-Asociación para la Prevención del Tabaquismo en Aragón  
SALT - Sociedad Aragonesa Libre de Tabaco  
Aireberri - Sociedad Vasconavarra para la Prevención del Tabaquismo  
APEPOC - Asociación de Pacientes con EPOC  
SOCIDROGALCOHOL  
FAECAP - Federación de Enfermería Comunitaria y de Atención Primaria  
Societat Catalana d'Atenció i Tractament del Consum del Tabac (SCATT)



## SWEDEN

A Non Smoking Generation  
Heart Lung Foundation  
Tobaksfacta  
Doctors against Tobacco  
Psychologists, Social Workers and Health Promoters against Tobacco  
Dentistry against Tobacco  
Nurses Association for a Tobacco Free Society  
Teachers against Tobacco



## TÜRKİYE

The Health Institute Association (Sağliga Evet Derneği).  
YESILAY - Turkish Green Crescent Society.  
Kansersiz Yaşam Derneği (KAYD).



## UKRAINE

Advocacy Center Life



## SWITZERLAND

The Swiss Tobacco Prevention Association  
OxySuisse



## UNITED STATES

Action on Smoking & Health -ASH  
Campaign for Tobacco-Free Kids

## ABOUT ENSP

The European Network for Smoking and Tobacco Prevention (ENSP) is an international organisation based in Brussels, active for 29 years in the field of tobacco control.

The organization was established between 1995 and 1997 at the request of the European Commission as part of the Europe against Cancer (EaC) Programme (1994 Vinci Resolution). Its creation aimed to bring together, under a unified European framework, key stakeholders engaged in tobacco control and cancer prevention. By consolidating expertise, advocacy, and policy engagement at the European level, the organization sought to strengthen coordination, amplify the impact of public health initiatives, and support the development of coherent, evidence-based strategies to reduce the burden of noncommunicable diseases across Europe[3].

With more than 80 members, including national coalitions, present in 38 European Countries (23 EU and 15 non-EU), ENSP objectives are to fully implement the WHO Framework Convention on Tobacco Control (FCTC) and to reduce the prevalence of tobacco use in the European region to less than 5% by 2040.

# ANNEX

## Evidence Summary: Preventing Nicotine Addiction Among Young People in Europe

The evidence demonstrates that nicotine is a highly addictive psychoactive substance with specific and heightened risks for adolescents and that current market developments are increasing exposure and dependence among young people in Europe.

Nicotine exerts its addictive effects primarily through activation of nicotinic acetylcholine receptors (nAChRs), leading to stimulation of the mesolimbic dopamine system and reinforcement of repeated use (Jiang et al., 2025). Chronic exposure produces neuroadaptations, including receptor desensitisation and upregulation, which sustain dependence (Liu et al., 2025). Importantly, adolescent exposure is associated with long-term alterations in brain development, including changes in reward processing and impulse control, increasing vulnerability to addiction and other substance use later in life (Reynolds et al., 2025; McGrath-Morrow et al., 2020). These findings confirm that nicotine addiction is not only a behavioural phenomenon but a neurobiological process with heightened impact during youth.

Recent product innovation has significantly increased the addiction potential of nicotine, particularly among young users. The introduction of nicotine salt formulations allows higher concentrations of nicotine to be delivered with reduced sensory irritation, facilitating rapid absorption and higher peak plasma levels (Benowitz, 2010; Jackler & Ramamurthi, 2019). Flavourings and product design features further enhance appeal and reduce aversion, especially among adolescents (Leventhal et al., 2021; Gades et al., 2022). Survey research found that use of sweet-flavoured Electronic Nicotine Delivery Systems (ENDS) increased over time with tobacco flavour decreasing in popularity, suggesting that flavoured products have greater abuse potential (Gades et al., 2022). These developments result in pharmacokinetic profiles that increasingly resemble or exceed those of combustible cigarettes, thereby amplifying dependence risk.

Epidemiological evidence indicates a fundamental shift in nicotine use patterns, with non-combustible products now serving as the primary entry point for youth. Adolescents are significantly more likely than adults to use e-cigarettes, with prevalence estimates indicating that youth are on average nine times more likely to vape (World Health Organization, 2024). In the WHO European Region, approximately 11.6% of adolescents aged 13–15 years use tobacco products and this figure likely underestimates total nicotine exposure due to increasing use of e-cigarettes and nicotine pouches (World Health Organization, 2024).

In several settings, e-cigarette use among adolescents exceeds that of conventional cigarette smoking, indicating a shift rather than a reduction in nicotine consumption.

Evidence of dependence among youth users is substantial. Surveys show high levels of loss of autonomy across nicotine products, including among exclusive e-cigarette users (Lin et al., 2022). Increasing reports of craving and early-morning use suggest strengthening dependence over time, particularly following the introduction of high-dose nicotine salt products (Jackson et al., 2025). European data further indicate that a significant proportion of adolescents self-identify as addicted, with a majority of quit attempts unsuccessful (Raad et al., 2025). These findings demonstrate that modern nicotine products are not mitigating addiction but are actively facilitating its development.

Patterns of dual and poly-use are increasingly common among young people, involving combinations of e-cigarettes, cigarettes and oral nicotine products. Such patterns are associated with sustained nicotine exposure and reduced likelihood of cessation, thereby increasing cumulative health risks (Hamoud et al., 2024; Zamarripa et al., 2025). This reinforces the conclusion that new nicotine products often complement rather than replace existing forms of use.

Nicotine exposure during adolescence is associated with direct and long-term health consequences. It disrupts neurodevelopmental processes and is linked to cognitive impairment and increased susceptibility to neuropsychiatric disorders (McGrath-Morrow et al., 2020; Reynolds et al., 2025). Nicotine also contributes to cardiovascular dysfunction through mechanisms including oxidative stress, endothelial damage and sympathetic activation (Münzel et al., 2025). These biological effects indicate that nicotine is not a benign substance and that early exposure may initiate long-term disease pathways.

From a population health perspective, early initiation of nicotine use is strongly associated with greater dependence severity, longer duration of use and higher cumulative lifetime exposure (Munzel et al., 2021). This trajectory is likely to translate into a substantial future burden of cardiovascular, metabolic and neuropsychiatric disease across Europe. The increasing prevalence of youth nicotine use, combined with more efficient delivery technologies and highly appealing product designs, suggests that without strengthened preventive measures, a new generation may face prolonged exposure and associated health risks.

In conclusion, the evidence clearly indicates that nicotine addiction among young people is being driven by a combination of biological vulnerability, product engineering and changing patterns of use. Strengthening efforts to prevent youth nicotine initiation and dependence therefore represents a critical opportunity to protect public health and reduce long-term economic burden within the European Union.

## **Main References:**

1. Jiang J, Li X, Hu AF et al. Nicotine and neuronal nicotinic acetylcholine receptors: unraveling the mechanisms of nicotine addiction. *Front Neurosci.* 2025;19:1670883. doi:10.3389/fnins.2025.1670883.

- 2.Liu S, Chen H, Zhu X et al. Structural insights into the progressive recovery of  $\alpha 7$  nicotinic acetylcholine receptor from nicotine-induced desensitization. *Sci Adv.* 2025;11(41):eadx4432. doi:10.1126/sciadv.adx4432.
- 3.Reynolds LM, Faure P, Barik J. Adolescent nicotine exposure and persistent neurocircuitry changes: unveiling lifelong psychiatric risks. *Mol Psychiatry.* 2025;30:5534–5545. doi:10.1038/s41380-025-03110-0.
- 4.McGrath-Morrow SA et al. The effects of nicotine on development. *J Physiol.* 2020. PMCID: PMC7049940.
- 5.Benowitz NL. Nicotine addiction. *N Engl J Med.* 2010;362:2295–2303. doi: 10.1056/NEJMra0809890
- 6.Jackler RK, Ramamurthi D. Nicotine arms race: JUUL and the high-nicotine product market. *Tob Control.* 2019;28:623–628. doi: 10.1136/tobaccocontrol-2018-054796
- 7.Leventhal AM, Madden DR, Peraza N, Schiff SJ, Lebovitz L, Whitted L, Barrington-Trimis J, Mason TB, Anderson MK, Tackett AP. Effect of Exposure to e-Cigarettes With Salt vs Free-Base Nicotine on the Appeal and Sensory Experience of Vaping: A Randomized Clinical Trial. *JAMA Netw Open.* 2021 Jan 4;4(1):e2032757. doi: 10.1001/jamanetworkopen.2020.32757. PMID: 33433597; PMCID: PMC7804919.
- 8.Gades, Mari S., et al. "The role of nicotine and flavor in the abuse potential and appeal of electronic cigarettes for adult current and former cigarette and electronic cigarette users: a systematic review." *Nicotine and Tobacco Research* 24.9 (2022): 1332-1343.
- 9.WHO. Global report on trends in prevalence of tobacco use 2000–2024 and projections 2025–2030. 2024. ISBN 978-92-4-011627-6. Available at: <https://www.who.int/publications/i/item/9789240116276>
- 10.Lin C, Gaiha SM, Halpern-Felsher B. Nicotine Dependence from Different E-Cigarette Devices and Combustible Cigarettes among US Adolescent and Young Adult Users. *Int J Environ Res Public Health.* 2022;19:5846.
- 11.Jackson SE, Brown J, Tattan-Birch H, Jarvis MJ. Changing patterns of nicotine product use and nicotine dependence among United States high-school students: The National Youth Tobacco Survey, 2014–2023. *Addiction.* 2025;120:2215–2222.
12. Raad DCJ et al. Frequent, dual and nighttime nicotine use among Dutch adolescents: findings from a school-based survey. *Eur J Pediatr.* 2025;184:816.
- 13.Hamoud J et al. A Systematic Review Investigating the Impact of Dual Use of E-Cigarettes and Conventional Cigarettes on Smoking Cessation. *ERJ Open Res.* 2024. doi:10.1183/23120541.00902-2024.

- 14.Zamarripa CA et al. A comprehensive review on oral nicotine pouches: Available scientific evidence and future research needs. *Exp Clin Psychopharmacol*. 2025;33:123–132.
- 15.Munzel T, Crea F, Rajagopalan S, Luscher T. Nicotine and the cardiovascular system: unmasking a global public health threat. *Eur Heart J*. 2025. doi: 10.1093/eurheartj/ehaf1010
- 16.Munzel T, Sorensen M, Daiber A. Transportation noise pollution and cardiovascular disease. *Nat Rev Cardiol*. 2021;18:619–636. doi: 10.1038/s41569-021-00532-5

## Extended list of relevant references

- 1.Al-Otaibi HM, Althobiani MA. Nicotine pouches: a narrative review of the existing literature. *Front Public Health*. 2025;13:1641308.
- 2.Alcheva A, Riegelman AL, Hatsukami DK. The Role of Nicotine and Flavor in the Abuse Potential and Appeal of Electronic Cigarettes for Adult Current and Former Cigarette and Electronic Cigarette Users: A Systematic Review. *Nicotine Tob Res*. 2022;24:1332–1343.
- 3.Ali M, Gray TR, Martinez DJ, Curry LE, Horn KA. Risk Profiles of Youth Single, Dual, and Poly Tobacco Users. *Nicotine Tob Res*. 2016;18:1614–1621.
- 4.Ali M, Gray TR, Martinez DJ, Curry LE, Horn KA. Risk profiles of youth single, dual, and poly tobacco users. *Nicotine Tob Res*. 2016;18(7):1614-1621. doi:10.1093/ntr/ntw028
- 5.Bals R, Boyd J, Esposito S, et al. Electronic cigarettes: a task force report from the European Respiratory Society. *Eur Respir J* 2019;53:1801151.
- 6.Banks E, Yazidjoglou A, Brown S, et al. Electronic cigarettes and health outcomes: Umbrella and systematic review of the global evidence. *Med J Aust*. 2023;218(6):267-275. doi: 10.5694/mja2.51890
- 7.Benowitz NL, Burbank AD. Cardiovascular toxicity of nicotine: Implications for electronic cigarette use. *Trends Cardiovasc Med*. 2016;26(6):515–23. doi:10.1016/j.tcm.2016.03.001.
- 8.Benowitz NL, Hukkanen J, Jacob P, 3rd. Nicotine chemistry, metabolism, kinetics and biomarkers. *Handb Exp Pharmacol*. 2009:29–60. doi: 10.1007/978-3-540-69248-5\_2
- 9.Benowitz NL, St Helen G, Liakoni E. Clinical Pharmacology of Electronic Nicotine Delivery Systems (ENDS): Implications for Benefits and Risks in the Promotion of the Combusted Tobacco Endgame. *J Clin Pharmacol*. 2021 Aug;61 Suppl 2(Suppl 2):S18-S36. doi: 10.1002/jcph.1915. PMID: 34396553; PMCID: PMC9239851.
- 10.Benowitz NL, Yang HY, Jacob P, St Helen G. Nicotine Dosimetry in Evaluating Electronic Cigarettes Compared to Cigarette Smoking: Implications for Tobacco Regulatory Science. *Chem Res Toxicol*. 2025;38:686–694.

11. Benowitz NL. Nicotine addiction. *N Engl J Med*. 2010;362:2295–2303. doi: 10.1056/NEJMra0809890
12. Benowitz NL. Nicotine and cardiovascular disease. *Prog Cardiovasc Dis*. 2016;58:146–156.
13. Benowitz NL. Pharmacology of nicotine: addiction, smoking-induced disease, and therapeutics. *Annu Rev Pharmacol Toxicol*. 2009;49:57–71. doi:10.1146/annurev.pharmtox.48.113006.094742.
14. Benowitz, N. L., St. Helen, G., & Liakoni, E. (2021). Clinical pharmacology of electronic nicotine delivery systems (ENDS): implications for benefits and risks in the promotion of the combusted tobacco endgame. *The Journal of Clinical Pharmacology*, 61, S18-S36.
15. Berman ML, Zettler PJ, Jordt SE. Synthetic Nicotine: Science, Global Legal Landscape, and Regulatory Considerations. *World Health Organ Tech Rep Ser*. 2023;1047:35–60. PMID: 37745838.
16. Berry KM, Fetterman JL, Benjamin EJ, et al. Association of electronic cigarette use with subsequent initiation of tobacco cigarettes in US youths. *JAMA Netw Open*. 2019;2(2):e187794. doi:10.1001/jamanetworkopen.2018.7794
17. Bittoni MA, Carbone DP, Harris RE. Vaping, smoking and lung cancer risk. *J Oncol Res Ther* 2024;9:10229.
18. Bjurlin MA, Matulewicz RS, Roberts TR, et al. Carcinogen biomarkers in the urine of electronic cigarette users and implications for the development of bladder cancer: a systematic review. *Eur Urol Oncol* 2021;4:766–83.
19. Bold KW, Kong G, Cavallo DA, Camenga DR, Krishnan-Sarin S. Reasons for trying e-cigarettes and risk of continued use. *Pediatrics*. 2016;138(3):e20160895. doi:10.1542/peds.2016-0895
20. Borowiecki M, Emery SL, Kostygina G. New recreational nicotine lozenges, tablets, gummies and gum proliferate on the US market. *Tob Control*. 2024;33:414–416.
21. Bowe AK, Doyle F, Stanistreet D, et al. E-cigarette-only and dual use among adolescents in Ireland: emerging behaviours with different risk profiles. *Int J Environ Res Public Health*. 2021;18(1):332. doi:10.3390/ijerph18010332
22. Brose L, Bunce L, Cheeseman H. Prevalence of Nicotine Pouch Use Among Youth and Adults in Great Britain—Analysis of Cross-Sectional, Nationally Representative Surveys. *Nicotine Tob Res*. 2026;28:213–222.
23. Brossard P, Weitkunat R, Poux V, Lama N, Haziza C, Picavet P, Baker G, Lüdicke F. Nicotine pharmacokinetic profiles of the Tobacco Heating System 2.2, cigarettes and nicotine gum in Japanese smokers. *Regul Toxicol Pharmacol*. 2017 Oct;89:193-199. doi: 10.1016/j.yrtph.2017.07.032. Epub 2017 Jul 29. PMID: 28760390.

24. Bruin JE, Gerstein HC, Holloway AC. Long-term consequences of fetal and neonatal nicotine exposure. *Toxicological Sciences*. 2010;116(2):364–374. <https://doi.org/10.1093/toxsci/kfq103>
25. Budney AJ, Sargent JD, Lee DC. Vaping cannabis (marijuana): parallel concerns to e-cigs? *Addiction*. 2015;110(11):1699-1704. doi:10.1111/add.13036
26. Bullen C, Howe C, Laugesen M, et al. Electronic cigarettes for smoking cessation: a randomised controlled trial. *Lancet*. 2013;382(9905):1629-1637. doi:10.1016/S0140-6736(13)61842-5
27. Butler AR, Lindson N, Livingstone-Banks J, Notley C, Turner T, Rigotti NA, Fanshawe TR, Dawkins L, Begh R, Wu AD, Brose L, Conde M, Simonavičius E, Hartmann-Boyce J. Interventions for quitting vaping. *Cochrane Database Syst Rev*. 2025 Jan 8;1(1):CD016058. doi: 10.1002/14651858.CD016058.pub2. Update in: *Cochrane Database Syst Rev*. 2024 May 21;5:CD016058. doi: 10.1002/14651858.CD016058. PMID: 39777614; PMCID: PMC11706636.
28. Cahill K, Stevens S, Perera R, Lancaster T. Pharmacological interventions for smoking cessation: an overview and network meta-analysis. *Cochrane Database Syst Rev*. 2013;2013(5):CD009329. doi:10.1002/14651858.CD009329.pub2.
29. Cai J, Bidulescu A. Associations between e-cigarette use or dual use of e-cigarette and combustible cigarette and metabolic syndrome: results from the National Health and Nutrition Examination Survey (NHANES). *Ann Epidemiol* 2023;85:93–9.e2.
30. Carlsson S et al. New nicotine and tobacco products and risk of major diseases: A review of the scientific evidence. *Scand J Public Health*. 2025. doi:10.1177/14034948251391690.
31. Castro EM, Lotfipour S, Leslie FM. Nicotine on the developing brain. *Pharmacol Res*. 2023;190:106716.
32. Centers for Disease Control and Prevention (CDC). E-cigarette Use Among Youth and Young Adults. Atlanta: CDC; 2024.
33. Centner AM, Bhide PG, Salazar G. Nicotine in Senescence and Atherosclerosis. *Cells*. 2020;9:1035. doi:10.3390/cells9041035.
34. Chand BR, Hosseinzadeh H. Association between e-cigarette use and asthma: a systematic review and meta-analysis. *J Asthma* 2022;59:1722–31.
35. Chang YP, Seo YS. E-cigarette use and concurrent risk behaviors among adolescents. *Nurs Outlook*. 2021;69(3):302-310. doi:10.1016/j.outlook.2020.09.005
36. Chaumont M, de Becker B, Zaher W, Culie A, Deprez G, Melot C, Reye F, Van Antwerpen P, Delporte C, Debbas N, et al. Differential Effects of E-Cigarette on Microvascular Endothelial Function, Arterial Stiffness and Oxidative Stress: A Randomized Crossover Trial. *Sci Rep*. 2018;8:10378. doi: 10.1038/s41598-018-28723-0

- 37.Chen C, Huo C, Matthey-Mora PP, et al. Assessing the association between e-cigarette use and cardiovascular disease: A meta-analysis of exclusive and dual use with combustible cigarettes. *Addict Behav* 2024;157:108086.
- 38.Cheng WL, Chen KY, Lee KY, Feng PH, Wu SM. Nicotinic-nAChR signaling mediates drug resistance in lung cancer. *J Cancer*. 2020;11:1125–1140. doi: 10.7150/jca.36359
- 39.Cho YJ et al. E-Cigarette Nicotine Delivery Among Young Adults by Nicotine Form, Concentration, and Flavor: A Crossover Randomized Clinical Trial. *JAMA Netw Open*. 2024;7:e2426702.
- 40.Choi K, Forster JL. Beliefs and experimentation with electronic cigarettes: a prospective analysis among young adults. *Am J Prev Med*. 2014;46(2):175-178. doi:10.1016/j.amepre.2013.10.007
- 41.Ciscato M, Chouvaeff M, Mourot A. Nicotinic Receptors in the Medial Habenula to Interpeduncular Nucleus Pathway: Modulators of Reward, Aversion and Emotion. *Eur J Neurosci*. 2025;62(11):e70352. doi:10.1111/ejn.70352.
- 42.Coleman BN, Apelberg BJ, Ambrose BK, et al. Association between electronic cigarette use and openness to cigarette smoking among US young adults. *Nicotine Tob Res*. 2015;17(2):212-218. doi:10.1093/ntr/ntu211
- 43.Dai HD, Leventhal AM. Use of Traditional Smokeless, Snus, and Dissolvable Tobacco Among U.S. Youth. *Am J Prev Med*. 2023;64(2):204–212. doi:10.1016/j.amepre.2022.09.011.
- 44.Dautzenberg B et al. Transdermal nicotine in non-smokers: A systematic review to design COVID-19 clinical trials. *Respiratory Medicine and Research*. 2021;80:100844.
- 45.Demissie Z, Everett Jones S, Clayton HB, King BA. Adolescent risk behaviors and use of electronic vapor products and cigarettes. *Pediatrics*. 2017;139(2):e20162921. doi:10.1542/peds.2016-2921
- 46.Dennison Himmelfarb CR, Benowitz NL, Blank MD, Bhatnagar A, Chase PJ, Davis EM, Fetterman JL, Keller-Hamilton B, Ogungbe O, Page RL, 2nd, et al. Impact of Smokeless Oral Nicotine Products on Cardiovascular Disease: Implications for Policy, Prevention, and Treatment: A Policy Statement From the American Heart Association. *Circulation*. 2025;151:e1–e21. doi: 10.1161/cir.0000000000001293
- 47.Deo SH, Jenkins NT, Padilla J, Parrish AR, Fadel PJ. Norepinephrine increases NADPH oxidase-derived superoxide in human peripheral blood mononuclear cells via  $\alpha$ -adrenergic receptors. *Am J Physiol Regul Integr Comp Physiol*. 2013;305:R1124–1132. doi: 10.1152/ajpregu.00347.2013
- 48.Dowd AN et al. A Cross-Sectional Survey on Oral Nicotine Pouches: Characterizing Use-Motives, Topography, Dependence Levels, and Adverse Events. *Nicotine Tob Res*. 2024;26:245–249.

- 49.Duell AK et al. Nicotine in tobacco product aerosols: forms and implications. *Tob Control*. 2020;29:656–662. doi:10.1136/tobaccocontrol-2019-055275.
- 50.Duell AK, Pankow JF, Peyton DH. Free-Base Nicotine Determination in Electronic Cigarette Liquids by <sup>1</sup>H NMR Spectroscopy. *Chem Res Toxicol*. 2018;31(6):431–434. doi:10.1021/acs.chemrestox.8b00097.
- 51.Dunbar MS, Tucker JS, Ewing BA, et al. Frequency of e-cigarette use, health status, and risk and protective health behaviors in adolescents. *J Addict Med*. 2017;11(1):55-62. doi:10.1097/ADM.0000000000000272
- 52.Durkin K, Williford DN, Turiano NA, et al. Associations between peer use, costs and benefits, self-efficacy, and adolescent e-cigarette use. *J Pediatr Psychol*. 2021;46(1):112-122. doi:10.1093/jpepsy/jsaa097
- 53.El-Amin S, Kinnunen JM, Rimpelä A. Adolescents' perceptions of harmfulness of tobacco and tobacco-like products in Finland. *Int J Environ Res Public Health*. 2022;19(3):1485. doi:10.3390/ijerph19031485
- 54.Eng L, Evans WK, Eisele M, Nishimura KK, Land SR, Rami-Porta R, Dresler CM, Asamura H, et al. The International Association for the Study of Lung Cancer Staging Project: the impact of smoking status on lung cancer staging in the 9th edition of the TNM classification. *JTO*. 2025;20(10):1441–58.
- 55.England LJ, Bunnell RE, Pechacek TF, Tong VT, McAfee TA. Nicotine and the developing human: a neglected element in the electronic cigarette debate. *American Journal of Preventive Medicine*. 2015;49(2):286–293. <https://doi.org/10.1016/j.amepre.2015.01.015>
- 56.Ermann LL, Klefbom L. Swedish tobacco policy: Key learnings to decrease smoking and challenges that still lie ahead. *Tobacco Prevention & Cessation*. 2024;10(December):65. doi:10.18332/tpc/196350.
- 57.ESPAD. ESPAD Report 2019 – Results from the European School Survey Project on Alcohol and Other Drugs. EMCDDA Joint Publications; 2020. Available at: <https://www.espad.org/espad-report-2019>
- 58.European Medicines Agency. Development of new medicinal products for treatment of smoking: scientific guideline. Available at: <https://www.ema.europa.eu/en/development-new-medicinal-products-treatment-smoking-scientific-guideline>
- 59.Fadus MC, Smith TT, Squeglia LM. The rise of e-cigarettes, pod mod devices, and JUUL among youth: factors influencing use, health implications, and downstream effects. *Drug Alcohol Depend*. 2019;201:85-93. doi:10.1016/j.drugalcdep.2019.04.011
- 60.Fant R, Owen L, Henningfield J. Nicotine replacement therapy. *Prim Care Clin Off Pract*. 1999;26:633–652.

- 61.FDA, Center for Tobacco Products. Results from the Annual National Youth Tobacco Survey (NYTS). 2025. Available at: <https://www.fda.gov/tobacco-products/youth-and-tobacco/results-annual-national-youth-tobacco-survey-nyts>
- 62.Felicione NJ et al. Oral Nicotine Pouches: Rising Popularity and State of the Science. Public Health Reports. 2026;141:55–62.
- 63.Feliu A, Martínez C, Fernández E. Lights and shadows for public health: a critical analysis of the tobacco legislation in Spain. Luces y sombras para la salud pública: análisis crítico de la legislación sobre el tabaco en España. Gac Sanit. 2022;36(1):48-52. doi:10.1016/j.gaceta.2021.07.001
- 64.Fite PJ, Cushing CC, Poquiz J, Frazer AL. Family influences on the use of e-cigarettes. Journal of Substance Use. 2018;23(4):396-401. doi:10.1080/14659891.2018.1436601
- 65.Flierl MA, Rittirsch D, Nadeau BA, Chen AJ, Sarma JV, Zetoune FS, McGuire SR, List RP, Day DE, Hoesel LM, et al. Phagocyte-derived catecholamines enhance acute inflammatory injury. Nature. 2007;449:721–725. doi: 10.1038/nature06185
- 66.Gades, Mari S., et al. "The role of nicotine and flavor in the abuse potential and appeal of electronic cigarettes for adult current and former cigarette and electronic cigarette users: a systematic review." Nicotine and Tobacco Research 24.9 (2022): 1332-1343.
- 67.Gaiha SM, Lin C, Lempert LK, Halpern-Felsher B. Use, marketing, and appeal of oral nicotine products among adolescents, young adults, and adults. Addict Behav. 2023;140:107632. doi:10.1016/j.addbeh.2023.107632.
- 68.Gallus, S., Lugo, A., Liu, X., Borroni, E., Clancy, L., Gorini, G., ... & TackSHS Project Investigators. (2022). Use and awareness of heated tobacco products in Europe. Journal of Epidemiology, 32(3), 139-144.
- 69.Gentzke AS et al. Tobacco Product Use Among Middle and High School Students — United States, 2020. MMWR Morb Mortal Wkly Rep. 2020;69:1881–1888.
- 70.Glantz, Stanton A. "Heated tobacco products: the example of IQOS." Tobacco control 27.Suppl 1 (2018): s1-s6.
- 71.Glasser AM, Collins L, Pearson JL, et al. Overview of electronic nicotine delivery systems: a systematic review. Am J Prev Med. 2017;52(2):e33-e66. doi:10.1016/j.amepre.2016.10.036
- 72.Goniewicz ML, Knysak J, Gawron M, et al. Levels of selected carcinogens and toxicants in vapour from electronic cigarettes. Tob Control. 2014;23(2):133-139. doi:10.1136/tobaccocontrol-2012-050859
- 73.Government of Canada . Canadian Tobacco and Nicotine Survey (CTNS): summary of results for 2022. Government of Canada. Updated September 19, 2023. Accessed August 27, 2025. <https://www.canada.ca/en/health-canada/services/canadian-tobacco-nicotin...>

- 74.Griendling KK, Minieri CA, Ollerenshaw JD, Alexander RW. Angiotensin II stimulates NADH and NADPH oxidase activity in cultured vascular smooth muscle cells. *Circ Res*. 1994;74:1141–1148. doi: 10.1161/01.res.74.6.1141
- 75.GTSS Academy. Current e-cigarette use [online database]. CDC Foundation. Available at: <https://www.gtssacademy.org/>
- 76.Gummin DD, Mowry JB, Beuhler MC, Spyker DA, Rivers LJ, Feldman R, Brown K, Pham NPT, Bronstein AC, DesLauriers C. 2022 Annual Report of the National Poison Data System® (NPDS) from America's Poison Centers®: 40th Annual Report. *Clin Toxicol (Phila)*. 2023 Oct;61(10):717-939. doi: 10.1080/15563650.2023.2268981. Epub 2023 Dec 12. PMID: 38084513.
- 77.Hajek P, Pittaccio K, Pesola F, Myers Smith K, Phillips-Waller A, Przulj D. Nicotine delivery and users' reactions to Juul compared with cigarettes and other e-cigarette products. *Addiction*. 2020 Jun;115(6):1141-1148. doi: 10.1111/add.14936. Epub 2020 Jan 29. PMID: 31994254; PMCID: PMC7318270.
- 78.Hammond D, Reid JL, Burkhalter R, Hong D. Trends in smoking and vaping among young people: findings from the ITC youth survey. University of Waterloo; 2023.
- 79.Hammond D, Reid JL, Cole AG, Leatherdale ST. Electronic cigarette use and smoking initiation among youth: a longitudinal cohort study. *CMAJ*. 2017;189(43):E1328-E1336. doi:10.1503/cmaj.161002
- 80.Hamoud J et al. A Systematic Review Investigating the Impact of Dual Use of E-Cigarettes and Conventional Cigarettes on Smoking Cessation. *ERJ Open Res*. 2024. doi:10.1183/23120541.00902-2024.
- 81.Han DH et al. Nicotine Pouch and E-Cigarette Use and Co-Use Among US Youths in 2023 and 2024. *JAMA Netw Open*. 2025;8:e256739.
- 82.Hanewinkel R, Hansen J. Konsum von Nikotinbeuteln im Kindes- und Jugendalter. *Laryngorhinootologie*. 2025;104(8):505–512. doi:10.1055/a-2481-5202.
- 83.Hanewinkel R, Isensee B. Risk factors for e-cigarette, conventional cigarette, and dual use in German adolescents: a cohort study. *Prev Med*. 2015;74:59-62. doi:10.1016/j.ypmed.2015.03.006
- 84.Hartendorp APT et al. Determination of Nicotine Protonation State in E-Liquids by Low-Resolution Benchtop NMR Spectroscopy. *Chem Res Toxicol*. 2024;37(8):1283–1289. doi:10.1021/acs.chemrestox.3c00417.
- 85.Hartmann-Boyce J, Chepkin SC, Ye W, Bullen C, Lancaster T. Nicotine replacement therapy versus control for smoking cessation. *Cochrane Database Syst Rev*. 2018;5(5):CD000146. doi:10.1002/14651858.CD000146.pub5.

86. Harvanko AM, Havel CM, Jacob P, Benowitz NL. Characterization of Nicotine Salts in 23 Electronic Cigarette Refill Liquids. *Nicotine Tob Res.* 2020;22(7):1239–1243. doi:10.1093/ntr/ntz232.
87. Hatsukami DK, Meier E, Lindgren BR, et al. A randomized clinical trial examining the effects of instructions for electronic cigarette use on smoking-related behaviors and biomarkers of exposure. *Nicotine Tob Res.* 2020;22(9):1524–1532.
88. Havermans A et al. Awareness, use and perceptions of cigarillos, heated tobacco products and nicotine pouches: a survey among Dutch adolescents and adults. *Drug Alcohol Depend.* 2021;229(Pt B):109136. doi:10.1016/j.drugalcdep.2021.109136.
89. Herzog C, Jones A, Evans I, et al. Cigarette smoking and e-cigarette use induce shared DNA methylation changes linked to carcinogenesis. *Cancer Res* 2024;84:1898–914.
90. Heshmati J et al. Nicotine pouch pharmacokinetics compared to smoked tobacco: A systematic review and meta-analysis. *Drug Alcohol Depend Rep.* 2025;17:100389.
91. Heshmati J et al. Vaping cessation interventions: a systematic review and meta-analysis. *Tob Control.* 2025. doi:10.1136/tc-2024-058798.
92. Honeycutt L, Huerne K, Miller A, et al. A systematic review of the effects of e-cigarette use on lung function. *NPJ Prim Care Respir Med* 2022;32:45.
93. Hua M, Talbot P. Potential health effects of electronic cigarettes: A systematic review of case reports. *Prev Med Rep.* 2016 Jun 10;4:169-78. doi: 10.1016/j.pmedr.2016.06.002. PMID: 27413679; PMCID: PMC4929082.
94. Hukkanen, J., Jacob, P., & Benowitz, N. L. (2005). Metabolism and disposition kinetics of nicotine. *Pharmacological Reviews*, 57(1), 79–115.
95. Hutzler C, Paschke M, Kruschinski S, Henkler F, Hahn J, Luch A. Chemical hazards present in liquids and vapors of electronic cigarettes. *Arch Toxicol.* 2014;88(7):1295-1308. doi: 10.1007/s00204-014-1294-7
96. Inaba Y, Chen J-A, Bergmann SR. Prediction of future cardiovascular outcomes by flow-mediated vasodilatation of the brachial artery: a meta-analysis. *International Journal of Cardiovascular Imaging.* 2010;26:631–640. doi: 10.1007/s10554-010-9616-1
97. Jackler RK, Ramamurthi D. Nicotine arms race: JUUL and the high-nicotine product market. *Tob Control.* 2019;28:623–628. doi: 10.1136/tobaccocontrol-2018-054796
98. Jackson SE, Brown J, Tattan-Birch H, Jarvis MJ. Changing patterns of nicotine product use and nicotine dependence among United States high-school students: The National Youth Tobacco Survey, 2014–2023. *Addiction.* 2025;120:2215–2222.

99. Jamal A, Park-Lee E et al. Tobacco product use among middle and high school students – National Youth Tobacco Survey, United States, 2024. *MMWR*. 2024;73(41):917–24. doi:10.15585/mmwr.mm7341a2.
100. Jehl J, Ciscato M, Vicq E et al. The interpeduncular nucleus blunts the rewarding effect of nicotine. *Neuron*. 2025;113(12):1898–1907. doi:10.1016/j.neuron.2025.03.035.
101. Jiang J, Li X, Hu AF et al. Nicotine and neuronal nicotinic acetylcholine receptors: unraveling the mechanisms of nicotine addiction. *Front Neurosci*. 2025;19:1670883. doi:10.3389/fnins.2025.1670883.
102. Johnson NL, Patten T, Ma M, De Biasi M, Wesson DW. Chemosensory Contributions of E-Cigarette Additives on Nicotine Use. *Front Neurosci*. 2022;16:893587. doi:10.3389/fnins.2022.893587.
103. Kaludercic N, Mialet-Perez J, Paolocci N, Parini A, Di Lisa F. Monoamine oxidases as sources of oxidants in the heart. *Journal of molecular and cellular cardiology*. 2014;73:34–42. doi: 10.1016/j.yjmcc.2013.12.032
104. Kaur J et al. Health impacts of electronic nicotine delivery systems. *BMJ Open*. 2025. PMID: PMC12516984.
105. Kennedy CD, van Schalkwyk MCI, McKee M, et al. The cardiovascular effects of electronic cigarettes: a systematic review of experimental studies. *Prevent Med* 2019;127:105770.
106. Kent JT, Mok G, Austin E. Nicotine Toxicity From Repeat Use of Nicotine Pouches. *Nicotine Tob Res*. 2025 Mar 24;27(4):767-768. doi: 10.1093/ntr/ntae111. PMID: 39087735; PMCID: PMC11931207.
107. Kim K, Picciotto MR. Nicotine addiction: More than just dopamine. *Curr Opin Neurobiol*. 2023;83:102797. doi:10.1016/j.conb.2023.102797.
108. Kim S, Oancea SC. Electronic cigarettes may not be a ‘safer alternative’ of conventional cigarettes during pregnancy: evidence from the nationally representative PRAMS data. *BMC Pregnancy Childbirth* 2020;20:557.
109. Kim, Jinyoung, et al. "Awareness, experience and prevalence of heated tobacco product, IQOS, among young Korean adults." *Tobacco control* 27.Suppl 1 (2018): s74-s77.
110. Kinnunen JM, Rimpelä AH, Lindfors PL, et al. Electronic cigarette use among 14- to 17-year-olds in Europe. *Eur J Public Health*. 2021;31(2):402-408. doi:10.1093/eurpub/ckaa145
111. Krishnan-Sarin S, Morean ME, Camenga DR, Cavallo DA, Kong G. E-cigarette use among high school and middle school adolescents in Connecticut. *Nicotine Tob Res*. 2015;17(7):810-818. doi:10.1093/ntr/ntu243

- 112.Kundu A, Feore A, Abu-Zarour N, Sanchez S, Sutton M, Sachdeva K, Seth S, Schwartz R, Chaiton M. Evidence update on the respiratory health effects of vaping e-cigarettes: A systematic review and meta-analysis. *Tob Induc Dis*. 2025 Nov 19;23. doi: 10.18332/tid/209954. PMID: 41268099; PMCID: PMC12628669.
- 113.Kundu A, Feore A, Sanchez S, et al. Cardiovascular health effects of vaping e-cigarettes: A systematic review and meta-analysis. *Heart*. 2025;111(13):599-608. doi: 10.1136/heartjnl-2024-325030
- 114.Kundu A, Sachdeva K, Feore A, et al. Evidence update on the cancer risk of vaping e-cigarettes: A systematic review. *Tob Induc Dis*. 2025;23:10.18332/tid/192934. doi: 10.18332/tid/192934
- 115.Kundu A, Sanchez S, Seth S, et al. Evidence update on e-cigarette dependence: A systematic review and meta-analysis. *Addict Behav*. 2025;163:108243. doi: 10.1016/j.addbeh.2024.108243
- 116.Kuntic M, Oelze M, Steven S, Kröller-Schön S, Stamm P, Kalinovic S, Frenis K, Vujacic-Mirski K, Bayo Jimenez MT, Kvandova M, et al. Short-term e-cigarette vapour exposure causes vascular oxidative stress and dysfunction: evidence for a close connection to brain damage and a key role of the phagocytic NADPH oxidase (NOX-2). *Eur Heart J*. 2020;41:2472–2483. doi: 10.1093/eurheartj/ehz772
- 117.Le Borgne T, Nguyen C, Vicq E et al. Nicotine engages a VTA-NAc feedback loop to inhibit amygdala-projecting dopamine neurons and induce anxiety-like behaviors. *Nat Commun*. 2025;16:6196. doi:10.1038/s41467-025-61180-8.
- 118.Lee J, Yao Z, Boakye E, et al. The impact of chronic electronic cigarette use on endothelial dysfunction measured by flow-mediated vasodilation: A systematic review and meta-analysis. *Tob Induc Dis* 2024;22:1–10.
- 119.Leventhal AM, Goldenson NI, Cho J, et al. Flavored e-cigarette use and progression of vaping in adolescents. *Pediatrics*. 2019;144(5):e20190789. doi:10.1542/peds.2019-0789
- 120.Leventhal AM, Madden DR, Peraza N, Schiff SJ, Lebovitz L, Whitted L, Barrington-Trimis J, Mason TB, Anderson MK, Tackett AP. Effect of Exposure to e-Cigarettes With Salt vs Free-Base Nicotine on the Appeal and Sensory Experience of Vaping: A Randomized Clinical Trial. *JAMA Netw Open*. 2021 Jan 4;4(1):e2032757. doi: 10.1001/jamanetworkopen.2020.32757. PMID: 33433597; PMCID: PMC7804919.
- 121.Li J, Liu S, Cao G, Sun Y, Chen W, Dong F, Xu J, Zhang C, Zhang W. Nicotine induces endothelial dysfunction and promotes atherosclerosis via GTPCH1. *J Cell Mol Med*. 2018;22:5406–5417. doi: 10.1111/jcmm.13812
- 122.Li X, Zhang Y, Zhang R, Chen F, Shao L, Zhang L. Association between e-cigarettes and asthma in adolescents: a systematic review and meta-analysis. *Am J Prev Med* 2022;62:953–60.

- 123.Lin C, Gaiha SM, Halpern-Felsher B. Nicotine Dependence from Different E-Cigarette Devices and Combustible Cigarettes among US Adolescent and Young Adult Users. *Int J Environ Res Public Health*. 2022;19:5846.
- 124.Lindson N, Chepkin SC, Ye W, Fanshawe TR, Bullen C, Hartmann-Boyce J. Different doses, durations and modes of delivery of nicotine replacement therapy for smoking cessation. *Cochrane Database Syst Rev*. 2019;Issue 4:CD013308. doi:10.1002/14651858.CD013308.
- 125.Ling PM, Hrywna M, Talbot EM, Lewis MJ. Tobacco-derived nicotine pouch brands and marketing messages on Internet and traditional media: content analysis. *JMIR Form Res*. 2023;7:e39146. doi:10.2196/39146.
- 126.Liu S, Chen H, Zhu X et al. Structural insights into the progressive recovery of  $\alpha 7$  nicotinic acetylcholine receptor from nicotine-induced desensitization. *Sci Adv*. 2025;11(41):eadx4432. doi:10.1126/sciadv.adx4432.
- 127.Liu Y, Filippidis FT. Tobacco market trends in 97 countries between 2007 and 2021. *Tob Induc Dis*. 2024;22:1–10.
- 128.Lunell E, Fagerström K, Hughes J, Pendrill R. Pharmacokinetic Comparison of a Novel Non-tobacco-Based Nicotine Pouch (ZYN) With Conventional, Tobacco-Based Swedish Snus and American Moist Snuff. *Nicotine Tob Res*. 2020 Oct 8;22(10):1757-1763. doi: 10.1093/ntr/ntaa068. PMID: 32319528.
- 129.Mallock-Ohnesorg N, Rabenstein A, Stoll Y, Gertzen M, Rieder B, Malke S, Burgmann N, Laux P, Pieper E, Schulz T, et al. Small pouches, but high nicotine doses-nicotine delivery and acute effects after use of tobacco-free nicotine pouches. *Front Pharmacol*. 2024;15:1392027. doi: 10.3389/fphar.2024.1392027
- 130.Mallock, Nadja, et al. "Levels of nicotine and tobacco-specific nitrosamines in oral nicotine pouches." *Tobacco Control* 33.2 (2024): 193-199.
- 131.Mamy L, Pesce S, Sanchez W, Aviron S, Bedos C, Berny P, Bertrand C, Betoulle S, Charles S, Chaumot A, Coeurdassier M, Coutellec MA, Crouzet O, Faburé J, Fritsch C, Gonzalez P, Hedde M, Leboulanger C, Margoum C, Mougin C, Munaron D, Nélieu S, Pelosi C, Rault M, Sucre E, Thomas M, Tournebize J, Leenhardt S. Impacts of neonicotinoids on biodiversity: a critical review. *Environ Sci Pollut Res Int*. 2025 Feb;32(6):2794-2829. doi: 10.1007/s11356-023-31032-3. Epub 2023 Dec 1. PMID: 38036909.
- 132.Marques P, Piqueras L, Sanz MJ. An updated overview of e-cigarette impact on human health. *Respir Res*. 2021;22(1):151. doi: 10.1186/s12931-021-01737-5
- 133.McBride JS, Altman DG, Klein M, White W. Green tobacco sickness. *Tob Control*. 1998 Autumn;7(3):294-8. doi: 10.1136/tc.7.3.294. PMID: 9825425; PMCID: PMC1763894.
- 134.McGrath-Morrow SA et al. The effects of nicotine on development. *J Physiol*. 2020. PMCID: PMC7049940.

- 135.Meernik C, Baker HM, Kowitt SD, Ranney LM, Goldstein AO.. Impact of non-menthol flavours in e-cigarettes on perceptions and use: an updated systematic review. *BMJ Open*. 2019;9(10):e031598. Published 2019 Oct 16. doi: 10.1136/bmjopen-2019-031598
- 136.Mehra VM, Keethakumar A, Bohr YM, Abdullah P, Tamim H. The association between alcohol, marijuana, illegal drug use and current use of E-cigarette among youth and young adults in Canada: results from Canadian Tobacco, Alcohol and Drugs Survey 2017. *BMC Public Health*. 2019;19(1):1208. doi:10.1186/s12889-019-7546-y
- 137.Mehrotra R, Malhotra R, Siddiqi K, Mishra D. Global Epidemiology of Smokeless Tobacco Use. In: *Smokeless Tobacco and its Effects on Periodontal Health*. Springer, Cham; 2026. doi:10.1007/978-3-032-16409-4\_1.
- 138.Middlekauff HR. Cardiovascular impact of electronic-cigarette use. *Trends Cardiovasc Med* 2020;30:133–40.
- 139.Mihov Y, Hurlemann R. Altered amygdala function in nicotine addiction: insights from human neuroimaging studies. *Neuropsychologia*. 2012;50(8):1719–1729.
- 140.Milicic S, Leatherdale ST. The associations between E-cigarettes and binge drinking, marijuana use, and energy drinks mixed with alcohol. *J Adolesc Health*. 2017;60(3):320-327. doi:10.1016/j.jadohealth.2016.10.011
- 141.Mishra A et al. Harmful effects of nicotine. *Indian J Med Paediatr Oncol*. 2015. PMID: PMC4363846.
- 142.Morgan B, Dai X, Flor LS et al. Morbidity and mortality attributable to tobacco exposure: an analysis from the Global Burden of Disease Study 2023. *Tobacco Induced Diseases*. 2025;23(1):A54.
- 143.Mueller, M., Fadai, T., Rauh, J. et al. Nicotine reduces discrimination between threat and safety in the hippocampus, nucleus accumbens and amygdala. *Transl Psychiatry* 14, 319 (2024). <https://doi.org/10.1038/s41398-024-03040-5>
- 144.Munzel T, Crea F, Rajagopalan S, Luscher T. Nicotine and the cardiovascular system: unmasking a global public health threat. *Eur Heart J*. 2025. doi: 10.1093/eurheartj/ehaf1010
- 145.Münzel T, Hahad O, Kuntic M, Keaney JF, Deanfield JE, Daiber A. Effects of tobacco cigarettes, e-cigarettes, and waterpipe smoking on endothelial function and clinical outcomes. *Eur Heart J*. 2020;41(41):4057-4070. doi: 10.1093/eurheartj/ehaa460
- 146.Munzel T, Sorensen M, Daiber A. Transportation noise pollution and cardiovascular disease. *Nat Rev Cardiol*. 2021;18:619–636. doi: 10.1038/s41569-021-00532-5
- 147.Münzel Tea. Nicotine Products And The Cardiovascular System: A Global Health Challenge Across All Delivery Forms. *Nat Rev Cardiol*. 2026;in press.

- 138.Nara S, Yamaguti Y, Tsuda I. Review: Nicotinic acetylcholine receptors to regulate important brain activity. *Cogn Neurodyn*. 2023;18(3):1–6. doi:10.1007/s11571-023-09975-7.
- 149.National Institute for Health and Care Excellence (NICE). Tobacco dependence treatment guidance. 2023.
- 150.Newhouse P, Kellar K, Aisen P et al. Nicotine treatment of mild cognitive impairment: a 6-month double-blind pilot clinical trial. *Neurology*. 2012;78(2):91–101. doi:10.1212/WNL.0b013e31823efcbb.
- 151.NHS England Digital. Smoking, Drinking and Drug Use among Young People in England, 2023. 2024. Available at: <https://digital.nhs.uk/data-and-information/publications/statistical/smoking-drinking-and-drug-use-among-young-people-in-england/2023>
- 152.Nicksic NE, Barnes AJ. Is susceptibility to e-cigarettes among youth associated with tobacco and other substance use behaviors one year later? Results from the PATH study. *Prev Med*. 2019;121:109-114. doi:10.1016/j.ypmed.2019.02.006
- 153.Notes from the field: Nicotine pouch use among middle and high school students—United States, 2023–2024. *Morbidity and Mortality Weekly Report*.
- 154.Ollila H, Tarasenko Y, Ciobanu A, Lebedeva E, Raitasalo K. Exclusive and dual use of electronic cigarettes among European youth in 32 countries with different regulatory landscapes. *Tob Control*. 2024;33:622-627. doi:10.1136/tc-2022-057749
- 155.Otsuka R, Watanabe H, Hirata K, Tokai K, Muro T, Yoshiyama M, Takeuchi K, Yoshikawa J. Acute effects of passive smoking on the coronary circulation in healthy young adults. *JAMA*. 2001;286:436–441. doi: 10.1001/jama.286.4.436
- 156.Panagopoulos VN, Bailey A, Kostopoulos GK, Ioannides AA. Changes in distinct brain systems identified with fMRI during smoking cessation treatment with varenicline: a review. *Psychopharmacology (Berl)*. 2024 Apr;241(4):653-685. doi: 10.1007/s00213-024-06556-2. Epub 2024 Mar 2. PMID: 38430396; PMCID: PMC10927841.
- 157.Park-Lee E, Jamal A, Cowan H, et al. Notes from the Field: E-Cigarette and Nicotine Pouch Use Among Middle and High School Students — United States, 2024. *MMWR Morb Mortal Wkly Rep* 2024;73:774–778. DOI: <http://dx.doi.org/10.15585/mmwr.mm7335a3>
- 158.Patel A, Cook S, Mattingly DT, et al. Longitudinal association between exclusive and dual use of cigarettes and electronic nicotine delivery systems and asthma among U.S. adolescents. *J Adolesc Health*. 2023;73(3):437-444. doi: 10.1016/j.jadohealth.2023.04.009
- 159.Patel M, Kierstead EC, Kreslake J, Schillo BA. Patterns of oral nicotine pouch use among U.S. adolescents and young adults. *Prev Med Rep*. 2023 May 12;34:102239. doi: 10.1016/j.pmedr.2023.102239. PMID: 37228836; PMCID: PMC10203764.

160. Patrick ME, Miech RA, Carlier C, O'Malley PM, Johnston LD, Schulenberg JE. Self-reported reasons for vaping among 8th, 10th, and 12th graders in the US: Nationally-representative results. *Drug Alcohol Depend*. 2016;165:275-278. doi:10.1016/j.drugalcdep.2016.05.017
161. Patten T, De Biasi M.. History repeats itself: Role of characterizing flavors on nicotine use and abuse. *Neuropharmacology*. 2020;177:108162.
162. Pereira, C. E. P., J. C. G. Lira, and B. O. Ferreira. "Analysis of the risks of using e-cigarettes in pregnancy: an integrative review." *Res Soc Dev* 11.10 (2022): e185111032797.
163. Polosa R, Morjaria JB, Prosperini U, et al. Health effects in COPD smokers who switch to electronic cigarettes: a retrospective-prospective 3-year follow-up. *Int J Chron Obstruct Pulmon Dis*. 2018;13:2533-2542. doi:10.2147/COPD.S161138
164. Primack BA, Soneji S, Stoolmiller M, Fine MJ, Sargent JD. Progression to traditional cigarette smoking after electronic cigarette use among US adolescents and young adults. *JAMA Pediatr*. 2015;169(11):1018-1023. doi:10.1001/jamapediatrics.2015.1742
165. Public Health Agency of Sweden. Use of tobacco and nicotine products among adolescents [online]. 2024. Available at: [https://www.folkhalsomyndigheten.se/the-public-health-agency-of-sweden/living-conditions-and-lifestyle/andtg/tobacco/use-of-tobacco-and-nicotine-products/?utm\\_](https://www.folkhalsomyndigheten.se/the-public-health-agency-of-sweden/living-conditions-and-lifestyle/andtg/tobacco/use-of-tobacco-and-nicotine-products/?utm_)
166. Qeadan F, Nicolson A, Barbeau WA, et al. The association between dual use of electronic nicotine products and illicit drugs with adverse cardiovascular and respiratory outcomes in a longitudinal analysis using the Population Assessment of Tobacco and Health (PATH) survey. *Drug Alcohol Depend* 2023;7:100166.
167. Raad DCJ et al. Frequent, dual and nighttime nicotine use among Dutch adolescents: findings from a school-based survey. *Eur J Pediatr*. 2025;184:816.
168. Rahman M, Irmiler M, Introna M, et al. Insight into the pulmonary molecular toxicity of heated tobacco products using human bronchial and alveolar mucosa models at air-liquid interface. *Sci Rep* 2022;12:16396.
169. Regan AK, Bombard JM, O'Hegarty MM, et al. Adverse birth outcomes associated with prepregnancy and prenatal electronic cigarette use. *Obstet Gynecol* 2021;138:85–94.
170. Regan AK, Pereira G. Patterns of combustible and electronic cigarette use during pregnancy and associated pregnancy outcomes. *Sci Rep* 2021;11:13508.
171. Reid JL, Burkhalter R, Ball J et al. Awareness and use of oral nicotine pouches among youth and young adults, 2022–2024: repeat cross-sectional surveys in Canada, England, the USA and New Zealand. *BMJ Public Health*. 2025;3:e003457. doi:10.1136/bmjph-2025-003457.

- 172.Reynolds LM, Faure P, Barik J. Adolescent nicotine exposure and persistent neurocircuitry changes: unveiling lifelong psychiatric risks. *Mol Psychiatry*. 2025;30:5534–5545. doi:10.1038/s41380-025-03110-0.
- 173.Rinaldi, Selina, et al. "Oral nicotine pouches with an aftertaste? Part 2: in vitro toxicity in human gingival fibroblasts." *Archives of Toxicology* 97.9 (2023): 2343-2356.
- 174.Robichaud MO, Seidenberg AB, Byron MJ. Tobacco companies introduce 'tobacco-free' nicotine pouches. *Tob Control*. 2020 Dec;29(e1):e145-e146. doi: 10.1136/tobaccocontrol-2019-055321. Epub 2019 Nov 21. PMID: 31753961; PMCID: PMC7239723.
- 175.Saddleson ML, Kozlowski LT, Giovino GA, et al. Risky behaviors, e-cigarette use and susceptibility of use among college students. *Drug Alcohol Depend*. 2015;149:25-30. doi:10.1016/j.drugalcdep.2015.01.001
- 176.Sankara Rao Muditana et al. Adverse Effects and Therapeutic Potential of Nicotine: A Review. *Advances in Pharmacology and Pharmacy*. 2025;13(3):448–457. doi:10.13189/app.2025.130315.
- 177.Sansone L, Milani F, Fabrizi R et al. Nicotine: From Discovery to Biological Effects. *Int J Mol Sci*. 2023;24(19):14570. doi:10.3390/ijms241914570.
- 178.Schneller LM, Felicione NJ, Hammond D, Goniewicz ML, O'Connor RJ. Tobacco-Free Oral Nicotine Product Use Among Youth in the U.S., 2019–2021. *AJPM Focus*. 2023;2:100061.
- 179.Schuller HM. Is cancer triggered by altered signalling of nicotinic acetylcholine receptors? *Nat Rev Cancer*. 2009;9:195–205. doi: 10.1038/nrc2590
- 180.Siddiqi K, Husain S, Vidyasagaran A, Readshaw A, Mishu MP, Sheikh A. Global burden of disease due to smokeless tobacco consumption in adults: an updated analysis of data from 127 countries. *BMC Med*. 2020;18(1):222. doi:10.1186/s12916-020-01677-9.
- 181.Siddiqi TJ, Rashid AM, Siddiqi AK, et al. Association of electronic cigarette exposure on cardiovascular health: a systematic review and meta-analysis. *Curr Probl Cardiol* 2023;48:101748.
- 182.Singh S, Pillai S, Chellappan S. Nicotinic acetylcholine receptor signaling in tumor growth and metastasis. *J Oncology*. 2011;2011:256743.
183. Sjödin E, Andersson J, Nordendahl M, et al Thirty-six-year trends (1986–2022) in cigarette smoking and snus use in northern Sweden: a cross-sectional study *BMJ Open* 2024;14:e088162. doi: 10.1136/bmjopen-2024-088162
- 184.Slotkin, T. (2004). Cholinergic systems in brain development and disruption by neurotoxins: nicotine, environmental tobacco smoke, organophosphates. *Toxicology and Applied Pharmacology*, 198(2), 132–151. <https://doi.org/10.1016/j.taap.2003.06.001>

- 185.Sompa SI, Zettergren A, Ekström S, et al. Predictors of electronic cigarette use and its association with respiratory health and obesity in young adulthood in Sweden; findings from the population-based birth cohort BAMSE. *Environ Res* 2022;208:112760.
- 186.Soneji S, Barrington-Trimis JL, Wills TA, et al. Association between initial use of e-cigarettes and subsequent cigarette smoking among adolescents and young adults: a systematic review and meta-analysis. *JAMA Pediatr*. 2017;171(8):788-797. doi:10.1001/jamapediatrics.2017.1488
- 187.Soni S, Verma L. Nicotine and Neurotransmitters: an Update. *Research Journal of Pharmacy and Technology*. 2024;17(6):2605–2. doi:10.52711/0974-360X.2024.00407.
- 188.Special Eurobarometer 539: Attitudes of Europeans towards tobacco and related products. Brussels; 2023.
- 189.Spindel ER, McEvoy CT. The Role of Nicotine in the Effects of Maternal Smoking during Pregnancy on Lung Development and Childhood Respiratory Disease. Implications for Dangers of E-Cigarettes. *Am J Respir Crit Care Med*. 2016 Mar 1;193(5):486-94. doi: 10.1164/rccm.201510-2013PP. PMID: 26756937; PMCID: PMC4824926.
- 190.St. Helen, G., Havel, C., Dempsey, D. A., Jacob III, P., & Benowitz, N. L. (2016). Nicotine delivery, retention and pharmacokinetics from various electronic cigarettes. *Addiction*, 111(3), 535-544.
- 191.Sukhato, K., Anothaisintawee, T., Chawla, N. et al. Comparative effects of electronic cigarettes and dual use of electronic and conventional cigarettes on adverse pregnancy outcomes: a systematic review and meta-analysis. *BMC Public Health* 25, 4359 (2025). <https://doi.org/10.1186/s12889-025-25765-8>
- 192.Sultan Y et al. Smoking-Related Disease Impact in the Eastern Mediterranean Region: A Comprehensive Assessment Using Global Burden of Disease Data. *Asian Pac J Cancer Prev*. 2024;25(2):495–505. doi:10.31557/APJCP.2024.25.2.495.
- 193.Sun T et al. Global prevalence of heated tobacco product use, 2015–22: systematic review and meta-analysis. *Addiction*. 2023;118(8):1430–1444. doi:10.1111/add.16199.
- 194.Tabuchi, T., Gallus, S., Shinozaki, T., Nakaya, T., Kunugita, N., & Colwell, B. (2018). Heat-not-burn tobacco product use in Japan: its prevalence, predictors and perceived symptoms from exposure to secondhand heat-not-burn tobacco aerosol. *Tobacco control*, 27(e1), e25-e33.
- 195.Tang MS, Wu XR, Lee HW, et al. Electronic-cigarette smoke induces lung adenocarcinoma and bladder urothelial hyperplasia in mice. *Proc Natl Acad Sci U S A* 2019;116:21,727–31.
- 196.Tarassenko Y, Ciobanu A, Fayokun R, Lebedeva E, Commar A, Mauer-Stender K. Electronic cigarette use among adolescents in 17 European study sites: findings from the Global Youth Tobacco Survey. *Eur J Public Health*. 2022;32(1):126-132. doi:10.1093/eurpub/ckab180

- 197.The Health Consequences of Smoking: Nicotine Addiction: A Report of the Surgeon General. DHHS Publ. No CDC. Available at: <https://stacks.cdc.gov/view/cdc/22014> (1988).
- 198.Thomas KH, Davies NM, Taylor AE, Taylor GMJ, Gunnell D, Martin RM, Douglas I. Risk of neuropsychiatric and cardiovascular adverse events following treatment with varenicline and nicotine replacement therapy in the UK Clinical Practice Research Datalink: a case-cross-over study. *Addiction*. 2021;116:1532–1545. doi: 10.1111/add.15338
- 199.Thompson DA, Lehmler HJ, Kolpin DW, Hladik ML, Vargo JD, Schilling KE, LeFevre GH, Peebles TL, Poch MC, LaDuca LE, Cwiertny DM, Field RW. A critical review on the potential impacts of neonicotinoid insecticide use: current knowledge of environmental fate, toxicity, and implications for human health. *Environ Sci Process Impacts*. 2020 Jun 24;22(6):1315-1346. doi: 10.1039/c9em00586b. PMID: 32267911; PMCID: PMC11755762.
- 200.Treur JL, Rozema AD, Mathijssen JJP, van Oers H, Vink JM. E-cigarette and waterpipe use in two adolescent cohorts: cross-sectional and longitudinal associations with conventional cigarette smoking. *Eur J Epidemiol*. 2018;33(3):323-334. doi:10.1007/s10654-017-0345-9
- 201.Tsai J, Walton K, Coleman BN, et al. Reasons for electronic cigarette use among middle and high school students - National Youth Tobacco Survey, United States, 2016. *MMWR Morb Mortal Wkly Rep*. 2018;67(6):196-200. doi:10.15585/mmwr.mm6706a5
- 202.Tudor TE, Lotrean LM. Opinions and practices regarding electronic cigarette use among middle school students from rural areas of Romania. *Int J Environ Res Public Health*. 2022;19(12):7372. doi:10.3390/ijerph19127372
- 203.U.S. Department of Health and Human Services. Chapter 4 Nicotine Addiction: Past and Present. In: *How Tobacco Smoke Causes Disease—The Biology and Behavioral Basis for Smoking-Attributable Disease: A Report of the Surgeon General*. Atlanta (GA): Centers for Disease Control and Prevention; 2010.
- 204.U.S. Department of Health and Human Services. The Health Consequences of Smoking—50 Years of Progress: A Report of the Surgeon General. Atlanta, GA: U.S. Department of Health and Human Services, Centers for Disease Control and Prevention, National Center for Chronic Disease Prevention and Health Promotion, Office on Smoking and Health; 2014. Available at: <https://www.ncbi.nlm.nih.gov/books/NBK294308/>
- 205.U.S. Food and Drug Administration (FDA). (2023). Tobacco product use and associated factors among middle and high school students—United States. *Morbidity and Mortality Weekly Report*.
- 206.Unger JB, Barker J, Cruz TB, Leventhal AM, Pentz MA. Lucy – novel flavored nicotine gum, lozenges, and pouches: Are they misleading consumers? *Subst Use Misuse*. 2022;57(8):1328–31. doi:10.1080/10826084.2022.2076881.
- 207.Upadhyay S, Rahman M, Johanson G. Heated tobacco products: insights into composition and toxicity. *Toxics* 2023;11:667.

- 208.Ussher, M., Fleming, J. & Brose, L. Vaping during pregnancy: a systematic review of health outcomes. *BMC Pregnancy Childbirth* 24, 435 (2024). <https://doi.org/10.1186/s12884-024-06633-6>
- 209.Vallée A, Eid M, Feki A, Ayoubi JM. Maternal vaping and pregnancy adverse outcomes: A systematic review and meta-analysis. *Women Birth*. 2025 Sep;38(5):101951. doi: 10.1016/j.wombi.2025.101951. Epub 2025 Jul 11. PMID: 40651293.
- 210.Vardavas CI, Girvalaki C et al. Profile of incidental exposures to e-cigarette liquids in Europe, 2018–2019. *Hum Exp Toxicol*. 2021;40(6):1045–1050. doi:10.1177/0960327120975828.
- 211.Vibjerg H. Youth and the Use of Nicotine Products. *Tobacco Prevention & Cessation*. 2023;9(Supplement):A89. doi:10.18332/tpc/162727.
- 212.Villanueva-Blasco VJ, Belda-Ferri L, Vázquez-Martínez A. A systematic review on risk factors and reasons for e-cigarette use in adolescents. *Tob Induc Dis*. 2025;23:1–25.
- 213.Wang Q, Ji X, Rahman I. Dysregulated metabolites serve as novel biomarkers for metabolic diseases caused by e-cigarette vaping and cigarette smoking. *Metabolites* 2021;11:345.
- 214.Wang, X.; Chen, Y.; Dong, J.; Ge, J.; Liu, X.; Liu, J. Neurobiology of Stress-Induced Nicotine Relapse. *Int. J. Mol. Sci*. 2024, 25, 1482. <https://doi.org/10.3390/ijms25031482>
- 215.Warren GW, Romano MA, Kudrimodti MR, Randal ME, McGarry RC, Singh AK, et al. Nicotinic modulation of therapeutic response in vitro and in vivo. *Int J Cancer*. 2012;131:2519–27.
- 216.Warren GW, Singh AK. Nicotine and lung cancer. *J Carcinogenesis*. 2013;12(1):1–8.
217. Whitehead AK, Erwin AP, Yue X. Nicotine and vascular dysfunction. *Acta Physiol (Oxf)*. 2021;231:e13631. doi: 10.1111/apha.13631
- 218.WHO Study Group on Tobacco Product Regulation. Report on the scientific basis of tobacco product regulation: eighth report. Available at: <https://www.who.int/publications/i/item/9789240022720>
- 219.WHO. Global report on trends in prevalence of tobacco use 2000–2024 and projections 2025–2030. 2024. ISBN 978-92-4-011627-6. Available at: <https://www.who.int/publications/i/item/9789240116276>
- 220.WHO. Tobacco Fact Sheet. 2024. Available at: <https://www.who.int/news-room/fact-sheets/detail/tobacco>
- 221.WHO. WHO report on the global tobacco epidemic, 2023: protect people from tobacco smoke. <https://www.who.int/publications/i/item/9789240077164>. 2023.

- 222.Wills TA, Soneji SS, Choi K, et al. E-cigarette use and respiratory disorders: an integrative review of converging evidence from epidemiological and laboratory studies. *Eur Respir J* 2021; 57:1–16.
- 223.Wills TA, Soneji SS, Choi K, Jaspers I, Tam EK. E-cigarette use and respiratory disorders: An integrative review of converging evidence from epidemiological and laboratory studies. *Eur Respir J*. 2021;57(1):1901815. doi: 10.1183/13993003.01815-2019
- 224.Winnicka L, Shenoy MA. EVALI and the pulmonary toxicity of electronic cigarettes: A review. *J Gen Intern Med*. 2020;35(7):2130-2135. doi: 10.1007/s11606-020-05813-2
- 225.World Health Organization. Tobacco and environmental health impacts, 2017. Available at: <https://www.who.int/publications/i/item/9789241512497>
- 226.World Health Organization. WHO clinical treatment guideline for tobacco cessation in adults. Geneva: WHO; 2024. Available at: <https://www.who.int/publications/i/item/9789240096431>
- 227.Xian S, Chen Y. E-cigarette users are associated with asthma disease: A meta-analysis. *Clin Respir J* 2021;15:457–66.
- 228.Yang Z, Berhane K, Leventhal AM, Liu M, Barrington-Trimis JL, Thomas DC. Modeling the longitudinal transitions of electronic cigarettes and conventional cigarettes with time-dependent covariates among adolescents. *Prev Med*. 2022;164:107294. doi:10.1016/j.ypmed.2022.107294
- 229.Yeboah J, Crouse JR, Hsu FC, Burke GL, Herrington DM. Brachial flow-mediated dilation predicts incident cardiovascular events in older adults: the Cardiovascular Health Study. *Circulation*. 2007;115:2390–2397. doi: 10.1161/CIRCULATIONAHA.106.678276
- 230.Yin-Hui Leong, Ahmad Shalihin Mohd Samin. E-Cigarettes: Risks, Research and Challenges. Springer Nature Singapore; 2025. doi:10.1007/978-981-96-5087-3.
- 231.Yingst, J. M., Foulds, J., Veldheer, S., Hrabovsky, S., Trushin, N., Eissenberg, T. T., ... & Hobkirk, A. L. (2019). Nicotine absorption during electronic cigarette use among regular users. *PLoS One*, 14(7), e0220300.
- 232.Zamarripa CA et al. A comprehensive review on oral nicotine pouches: Available scientific evidence and future research needs. *Exp Clin Psychopharmacol*. 2025;33:123–132.
- 233.Zhang D, Zheng H, Cui K et al. Nicotinic acetylcholine receptors-targeting drug discovery. *Eur J Med Chem*. 2026;308:118679. doi:10.1016/j.ejmech.2026.118679.
- 234.Zhang Z, Jiao Z, Blaha MJ, et al. The association between e-cigarette use and prediabetes: results from the Behavioral Risk Factor Surveillance System, 2016–2018. *Am J Prev Med* 2022;62:872–7.